

## Theranostic effect of serial manganese-enhanced magnetic resonance imaging of human embryonic stem cell derived teratoma.

**Journal:** Magn Reson Med

**Publication Year:** 2011

**Authors:** J Chung, R Dash, K Kee, J K Barral, H Kosuge, R C Robbins, D Nishimura, Pera R Reijo, P C Yang

**PubMed link:** 22190225

**Funding Grants:** In Vivo Molecular Magnetic Resonance Imaging of Human Embryonic Stem Cells in Murine Model of Myocardial Infarction, The Stanford University Center for Human Embryonic Stem Cell Research and Education, Stanford CIRM Training Program

### Public Summary:

Although human embryonic stem cells (hESCs) hold therapeutic potential, teratoma formation has deterred clinical applications. Manganese ( $Mn^{2+}$ ) enters metabolically active cells and can generate high signal intensity on the T1 weighted magnetic resonance imaging. We hypothesized that serial manganese-enhanced MRI could assess hESC-derived teratoma formation and eliminate the teratoma by inducing cytotoxicity through intracellular accumulation of  $Mn^{2+}$ . hESCs were transplanted into severe combined immunodeficient mouse (SCID) hindlimbs to form teratoma. Then, these mice were divided into the two groups: chemotherapy group and control group. The chemotherapy group was injected with  $MnCl_2$  intraperitoneally three times a week. The control group was given  $MnCl_2$  only prior to MRI. Longitudinal evaluation by manganese-enhanced MRI was performed till 8 weeks post-transplant. The chemotherapy group showed significant reduction in the teratoma volume at weeks 6 and 8. Histology revealed increased proportion of dead cells and caspase 3 (apoptosis marker) positive cells in the chemotherapy group. In conclusion, systemic administration of  $MnCl_2$  enabled simultaneous monitoring and elimination of hESC-derived teratoma cells by higher intracellular accumulation of  $Mn^{2+}$ .

### Scientific Abstract:

Although human embryonic stem cell (hESC) hold therapeutic potential, teratoma formation has deterred clinical translation. Manganese ( $Mn^{2+}$ ) enters metabolically active cells through voltage-gated calcium channels and subsequently, induces T1 shortening. We hypothesized that serial manganese-enhanced MRI would have theranostic effect to assess hESC survival, teratoma formation, and hESC-derived teratoma reduction through intracellular accumulation of  $Mn^{2+}$ . Firefly luciferase transduced hESCs (hESC-Lucs) were transplanted into severe combined immunodeficient mouse hindlimbs to form teratoma. The chemotherapy group was injected with  $MnCl_2$  intraperitoneally three times a week. The control group was given  $MnCl_2$  only prior to manganese-enhanced MRI. Longitudinal evaluation by manganese-enhanced MRI and bioluminescence imaging was performed. The chemotherapy group showed significant reduction in the teratoma volume and luciferase activity at weeks 6 and 8. Histology revealed increased proportion of dead cells and caspase 3 positive cells in the chemotherapy group. Systemic administration of  $MnCl_2$  enabled simultaneous monitoring and elimination of hESC-derived teratoma cells by higher intracellular accumulation of  $Mn^{2+}$ . Magn Reson Med, 2011. (c) 2011 Wiley Periodicals, Inc.

**Source URL:** <http://www.cirm.ca.gov/about-cirm/publications/theranostic-effect-serial-manganese-enhanced-magnetic-resonance-imaging>